arXiv:1401.1652v2 [math.AG] 22 Dec 2015

ON CLASSIFICATION OF GROUPS OF POINTS ON ABELIAN VARIETIES OVER FINITE FIELDS

SERGEY RYBAKOV

To M. A. Tsfasman and S. G. Vlăduţ on the occasion of their 60th birthdays

Abstract. In this paper we improve our previous results on classification of groups of points on abelian varieties over finite fields. The classification is given in terms of the Weil polynomial of abelian varieties in a given k-isogeny class over a finite field k.

1. Introduction

Let X be an algebraic variety over a finite field $k = \mathbb{F}_q$ of characteristic p. The set $X(k)$ of points defined over k is an important invariant of X. If X is an abelian variety, then $X(k)$ is a finite abelian group. Previously [Ry12] we classified groups of points on abelian surfaces over finite fields. It is important that for the classification in question one first divide the set of abelian surfaces into isogeny classes, and then classify groups of points inside a given isogeny class. In this paper we follow the same strategy. By the Tate-Honda theorem, an isogeny class of abelian varieties over k corresponds to characteristic polynomial of Frobenius action on the ℓ-th Tate module of any variety from the class (the Weil polynomial). These polynomials are known for abelian varieties of low dimensions. In this paper we state some partial results concerning higher dimensions. In particular, we prove a conjecture stated in [Ry10]. We also clarify the proof of the classification theorem for groups of points on abelian surfaces.

2. Preliminary results on abelian varieties and notation

Let $\bar{k}$ be an algebraic closure of k. For a given prime number ℓ and a natural number m denote by A_m the kernel of multiplication by ℓ^m in $A(\bar{k})$. The ℓ-th Tate module of A is defined as $T_\ell(A) = \varprojlim A_m$. The Frobenius endomorphism F of A acts on the Tate module by a semisimple linear transformation, which we also denote by F. Suppose $\ell \neq p$. Then $T_\ell(A)$ is a free $\mathbb{Z}_\ell$-module of rank $2 \dim A$. We define *the Weil polynomial of* A as the characteristic polynomial

$$f_A(t) = \det(t - F|T_\ell(A)).$$

It is a monic polynomial of degree $2 \dim A$ with rational integer coefficients independent of the choice of prime ℓ. Tate proved that abelian varieties A and B are isogenous if and only if $f_A(t) = f_B(t)$ [Ta66].

In what follows we would like to treat the case $\ell = p$ as well. The module $T_p(A)$ sometimes is called the physical Tate module. The polynomial $g(t) = \det(t - F|T_p(A))$ divides $f_A(t)$ in $\mathbb{Z}_p[t]$. Moreover, $\deg g \leq \dim A$, and

$$f_A(t) \equiv g(t)t^{2 \dim A - \deg g} \mod p.$$

1991 *Mathematics Subject Classification.* 14K99, 14G05, 14G15.

Key words and phrases. abelian variety, the group of rational points, finite field, Newton polygon, Hodge polygon.

Supported in part by the research project DST-1211005, RFBR grants no. 12-01-31280, 12-01-92697-IND-a, 14-01-93108 and by a subsidy granted to the HSE by the Government of the Russian Federation for the implementation of the Global Competitiveness Program.

In fact, main results of this paper are trivial for T_p, because its rank is too small for abelian varieties of low dimension. However we think it is more natural to state the results for all primes.

Recall some notation and results from [Ry14, Section 2] . Let Δ be the set of roots of $f \bmod \ell$, and let $\Lambda \subset \Delta$ be the image of a section of the natural map $\Delta \to \Delta/\operatorname{Gal}(\overline{\mathbb{F}}_\ell/\mathbb{F}_\ell)$. Then $f = \prod_{\lambda\in\Lambda} f_\lambda$, where $f_\lambda \in \mathbb{Z}_\ell[t]$ is the monic polynomial such that the set of roots of $f_\lambda \bmod \ell$ is $\operatorname{Gal}(\overline{\mathbb{F}}_\ell/\mathbb{F}_\ell)\lambda$. We have a corresponding decomposition of the Tate module

$$T_\ell(A) \cong \oplus_{\lambda\in\Lambda} T_\lambda(A),$$

where F acts on $T_\lambda(A)$ with the characteristic polynomial f_λ. In this paper we are interested in the summand corresponding to $\lambda = 1$ which we denote by $T_\ell^{(1)}(A)$. We also write

$$T_\ell^{(2)}(A) = \oplus_{\lambda\neq 1} T_\lambda(A).$$

We denote by $f_\ell \in \mathbb{Z}_\ell[t]$ the polynomial f_1 corresponding to $\lambda = 1$. This is the divisor of f_A of maximal degree such that $f_\ell(t-1) \equiv t^{\deg f_\ell} \bmod \ell$. Note that this results hold for $\ell = p$ without any changes.

Let $f_\ell^{sep} \in \mathbb{Z}_\ell[t]$ be the monic separable polynomial with the same set of roots as f_ℓ, and let

$$R = \mathbb{Z}_\ell[t]/f_\ell^{sep}\mathbb{Z}_\ell[t].$$

Denote by $x \in R$ the image of t under the natural projection from $\mathbb{Z}_\ell[t]$. Thus $T_\ell^{(1)}(A)$ is an R-module such that x acts as F. Without danger of confusion we define *a Tate module* as an R-module which is free of finite rank over $\mathbb{Z}_\ell$. The *rank* of a Tate module is its rank over $\mathbb{Z}_\ell$.

For an abelian group G we denote by G_ℓ the ℓ-primary component of G. The group $A(k)$ is the kernel of $1 - F : A \to A$, and the ℓ-component

$$A(k)_\ell \cong T_\ell(A)/(1-F)T_\ell(A)$$

(see [Ry10, Proposition 3.1]). By definition, $T_\ell(A)/(1-F)T_\ell(A) \cong T_\ell^{(1)}(A)/(1-x)T_\ell^{(1)}(A)$. Informally, $T_\ell^{(1)}(A)$ is the minimal part of $T_\ell(A)$ which knows about the group of points. The proof of the following proposition is essentially the proof of [Ry10, Theorem 1.1].

Proposition 2.1. *Let f be a Weil polynomial, and let G be a finite abelian group of order $f(1)$. Suppose that for any prime number ℓ dividing $f(1)$ there exists a Tate module T_ℓ such that $f_\ell(t) = \det(t - x|T_\ell)$, and $G_\ell \cong T_\ell/(1-x)T_\ell$. Then there exists an abelian variety A over k such that $A(k) \cong G$.*

Proof. Let B be an abelian variety over k with Weil polynomial f. By the discussion before the proposition, $T_\ell(B) \cong T_\ell^{(1)}(B) \oplus T_\ell^{(2)}(B)$. Since the Frobenius action on the vector space $V_\ell^{(1)}(B) = T_\ell^{(1)}(B) \otimes \mathbb{Q}_\ell$ is semisimple, it is determined up to isomorphism by the polynomial f_ℓ. Therefore there exists an isomorphism $V_\ell^{(1)}(B) \cong T_\ell \otimes_{\mathbb{Z}_\ell} \mathbb{Q}_\ell$, and an inclusion $T_\ell \to V_\ell^{(1)}(B)$ such that the image of T_ℓ is contained in $T_\ell^{(1)}(B)$. Put $T'_\ell = T_\ell \oplus T_\ell^{(2)}(B)$. Clearly, we have an isomorphism of F-vector spaces $T'_\ell \otimes \mathbb{Q}_\ell \to V_\ell(B)$. By [Ry10, Lemma 2.1], there exists an abelian variety B_1 and an ℓ-isogeny $B_1 \to B$ such that $T_\ell(B_1) \cong T'_\ell$. By [Ry10, Proposition 3.1], we have $G_\ell \cong B_1(k)_\ell$.

Let $\ell_1, \dots, \ell_s$ be the set of prime divisors of $f(1)$. By the above there is a sequence of isogenies

$$A = B_s \xrightarrow{\varphi_s} B_{s-1} \to \dots \xrightarrow{\varphi_2} B_1 \xrightarrow{\varphi_1} B$$

such that $\varphi_i : B_i \to B_{i-1}$ is an ℓ_i-isogeny and

$$B_i(k)_{\ell_i} \cong G_{\ell_i}.$$

Since φ_i is an ℓ_i-isogeny, $T_\ell(B_i) \cong T_\ell(B_{i-1})$ for any $\ell \neq \ell_i$. Thus $A(k) \cong G$. □

Fix and a positive integer r. A *matrix factorization* (with respect to f_ℓ, f_ℓ^{sep}, and r) is a pair (X, Y) of $r \times r$ matrices with coefficients in $\mathbb{Z}_\ell[t]$ such that $\det X = f_\ell$ and $YX = f_\ell^{sep} \cdot I_r$, where I_r is the identity matrix. The following proposition shows the connection between Tate modules and matrix factorizations.

Proposition 2.2. [Ry12] *Let (X, Y) be a matrix factorization. Then*

$$T = \operatorname{coker}(\mathbb{Z}_\ell[t]^r \xrightarrow{X} \mathbb{Z}_\ell[t]^r). \tag{2.1}$$

is a Tate module, and the characteristic polynomial of the action of x on T is equal to f_ℓ.

Conversely, let T be a Tate module, which can be generated over R by r elements. Suppose that $\det(t-x|T) = f_\ell$, and $\operatorname{Hp}_\ell(T/(1-x)T, r) = (m_1, \dots, m_r)$. Then there exists a matrix factorization (X, Y) such that T has the presentation (2.1)*, and*

$$X \equiv \operatorname{diag}(\ell^{m_1}, \dots, \ell^{m_r}) \mod (1-t)\mathbb{Z}_\ell[t].$$

3. Tate modules and Hodge polygons

Denote by ν_ℓ the ℓ-adic valuation on $\mathbb{Q}_\ell$. Let $Q(t) = \sum_i Q_i t^i$ be a polynomial of degree d over $\mathbb{Q}_\ell$, and let $Q(0) \neq 0$. Take the lower convex hull of the points $(i, \nu_\ell(Q_i))$ for $0 \leq i \leq d$ in $\mathbb{R}^2$. The boundary of this region without vertical lines is called *the Newton polygon* $\operatorname{Np}_\ell(Q)$ *of* Q.

Let $G = \oplus_{i=1}^r \mathbb{Z}/\ell^{m_i}\mathbb{Z}$ be an ℓ–group of order ℓ^m. In what follows we assume that the exponents $m_1, \dots, m_r$ are ordered in the following way: $0 \leq m_1 \leq m_2 \leq \cdots \leq m_r$. Note that some of these numbers could be zero. The *Hodge polygon* $\operatorname{Hp}_\ell(G, r)$ *of the group* G is the convex polygon with vertices $(i, \sum_{j=1}^{r-i} m_j)$ for $0 \leq i \leq r$. It has $(0, m)$ and $(r, 0)$ as its endpoints, and its slopes are $-m_r, \dots, -m_1$. We write $\operatorname{Hp}_\ell(G, r) = (m_1, \dots, m_r)$. If T is a Tate module, we denote $\operatorname{Hp}_\ell(T/(1-x)T, \operatorname{rk} T)$ by $\operatorname{Hp}_\ell(T)$.

Lemma 3.1. *Let*

$$0 \to H \to G \to G' \to 0$$

be a short exact sequence of finite abelian ℓ–groups. Let $\operatorname{Hp}_\ell(H, s) = (n_1, \dots, n_s)$, and $\operatorname{Hp}_\ell(G, r) = (m_1, \dots, m_r)$. Assume that G' is generated by $r - s$ elements. Then $m_i \leq n_i$ for $1 \leq i \leq s$.

Dually, if $\operatorname{Hp}_\ell(G', r-s) = (n'_1, \dots, n'_{r-s})$, then $m_i \geq n'_i$ for $s+1 \leq i \leq r$.

Proof. The group G is isomorphic to

$$\oplus_{i=1}^r \mathbb{Z}/\ell^{m_i}\mathbb{Z}.$$

Choose $v_1, \dots, v_r \in G$ such that v_i generates the summand $\mathbb{Z}/\ell^{m_i}\mathbb{Z}$. We choose generators $u_1, \dots, u_s \in H$ in the same way. Let H' be a subgroup of H generated by $u_1, \dots, u_i$. The group G' is generated by $r-s$ elements; thus G/H' is generated by $r-i$ elements, and there exists a linear combination

$$v = \sum_{j=i}^{r} a_j v_j$$

such that $v \in H'$, and there exists j such that $\nu_\ell(a_j) = 0$. Now assume that $m_i > n_i$. It follows that $\ell^{n_i} v \neq 0$. On the other hand, $\ell^{n_i} u = 0$ for any $u \in H'$. This contradiction concludes the proof.

The last assertion can be proved by the dual argument. □

Theorem 3.2. *Let T be a Tate module of rank r, and T_1 be a Tate submodule of rank s such that $T_2 = T/T_1$ is torsion-free. Suppose that $\operatorname{Hp}_\ell(T_1) = (n_1, \dots, n_s)$, and $\operatorname{Hp}_\ell(T) = (m_1, \dots, m_r)$. Then $m_i \leq n_i$ for $1 \leq i \leq s$.*

Dually, if $\operatorname{Hp}_\ell(T_2) = (n'_1, \dots, n'_{r-s})$, then $m_i \geq n'_i$ for $s+1 \leq i \leq r$.

Proof. We prove that $m_i \le n_i$ for $1 \le i \le s$. The second assertion follows from the dual argument. The transformation $E = 1 - x$ acts on the short exact sequence of Tate modules:

$$0 \to T_1 \to T \to T_2 \to 0.$$

By the snake lemma we get a exact sequence of abelian groups:

$$0 \to T_1/ET_1 \to T/ET \to T_2/ET_2 \to 0.$$

The Lemma 3.1 completes the proof. □

Lemma 3.3. *Let L be a finite extension of $\mathbb{Q}_\ell$ generated by an element α with ℓ-adic valuation $\lambda \in \mathbb{Q}$. Denote by $\mathcal{O}$ the ring of integers of L. Then there exists an isomorphism of abelian groups:*

$$\mathcal{O}/\alpha\mathcal{O} \cong (\mathbb{Z}/\ell^{\lfloor\lambda\rfloor}\mathbb{Z})^{[L:\mathbb{Q}_\ell]-e} \oplus (\mathbb{Z}/\ell^{\lceil\lambda\rceil}\mathbb{Z})^e,$$

where $\lambda = \lfloor\lambda\rfloor + e/[L:\mathbb{Q}_\ell]$.

The proof is an easy application of [La94, Proposition I.7.23].

Theorem 3.4. *Let T be a Tate module of rank r, and let $\mathrm{Hp}_\ell(T) = (m_1, \dots, m_r)$. Put $Q(t) = f(1-t)$. Let α be a root of Q with multiplicity s and ℓ-adic valuation*

$$\lambda = \lfloor\lambda\rfloor + \frac{e}{[L:\mathbb{Q}_\ell]}, \text{ and let } s' = \left\lceil \frac{([L:\mathbb{Q}_\ell]-e)s}{[L:\mathbb{Q}_\ell]} \right\rceil,$$

where $L = \mathbb{Q}_\ell(\alpha)$. Then

$$m_{s'} \le \lfloor\lambda\rfloor,\ m_s \le \lceil\lambda\rceil,\ m_{r-s'+1} \ge \lceil\lambda\rceil, \text{ and } m_{r-s+1} \ge \lfloor\lambda\rfloor.$$

Proof. Denote by $\mathcal{O}$ the ring of integers of L. Let $S = T \otimes_{\mathbb{Z}_\ell} \mathcal{O}$. The Tate module S contains the submodule $S_1 = \ker(1 - x - \alpha)$. Clearly,

$$S_1/(1-x)S_1 \cong (\mathcal{O}/\alpha\mathcal{O})^s, \text{ and } S/(1-x)S \cong (T/(1-x)T) \otimes_{\mathbb{Z}_\ell} \mathcal{O} \cong (T/(1-x)T)^{[L:\mathbb{Q}_\ell]}.$$

By Lemma 3.3, the Hodge polygon of $(\mathcal{O}/\alpha\mathcal{O})^s$ has $([L:\mathbb{Q}_\ell]-e)s$ slopes $\lfloor\lambda\rfloor$ and es slopes $\lceil\lambda\rceil$. Thus the slope number $[L:\mathbb{Q}_\ell]s$ is equal to $\lceil\lambda\rceil$. By Theorem 3.2, $m_{s'} \le \lfloor\lambda\rfloor$, and $m_s \le \lceil\lambda\rceil$. The rest can be proved dually. □

Corollary 3.5. *Let A be an abelian variety over k, and let $\mathrm{Hp}_\ell(T_\ell^{(1)}(A)) = (n_1, \dots, n_r)$. Put $T = \wedge^e T_\ell(A)$, and $\mathrm{Hp}_\ell(T) = (m_1, \dots, m_{r'})$, where $r' = \binom{r}{e}$. Clearly, for any i there exist $i_1 < \dots < i_e$ such that $m_i = n_{i_1} + \dots + n_{i_e}$. Let $Q(t) = \det(t - E|T)$, where E is induced by $1 - x$. Let α be a root of Q with multiplicity s and ℓ-adic valuation $\lambda \in \mathbb{Q}$. Then $m_s \le \lceil\lambda\rceil$, and $m_{r'-s+1} \ge \lfloor\lambda\rfloor$.*

4. Main results

Theorem 4.1. *Let $f = P^r Q^s$ be a Weil polynomial, where Q divides P, and $\deg P \le 2$. Suppose that P is separable. Then G is the group of points on some abelian variety with Weil polynomial f if and only if*

$$G_\ell \cong \oplus_{j=1}^r G_\ell^{(j)} \oplus H,$$

where $G_\ell^{(j)}$ are ℓ-primary abelian groups such that $\mathrm{Np}_\ell(P(1-t))$ lies on or above $\mathrm{Hp}_\ell(G_\ell^{(j)}, 2)$ for all $1 \le j \le r$, and $H \cong (\mathbb{Z}_\ell/Q(1)\mathbb{Z}_\ell)^s$.

Proof. If $\deg P_\ell \le 1$, the statement is trivial. We assume that $P_\ell = P$, and $\deg P = 2$; therefore $f_\ell = f$. It is easy to construct a variety with a given group of points. By Proposition 2.1, we have to construct Tate modules T_ℓ over $R = \mathbb{Z}_\ell[t]/P(t)\mathbb{Z}_\ell[t]$ such that $T_\ell/(1-x)T_\ell \cong G_\ell$. By [Ry10, Theorem 3.2], there exist Tate modules $T_\ell^{(j)}$ such that

$$T_\ell^{(j)}/(1-x)T_\ell^{(j)} \cong G_\ell^{(j)}$$

and P is the characteristic polynomial of Frobenius action on $T_\ell^{(j)}$. Clearly, there exists a Tate module T_H such that $T_H/(1-x)T_H \cong H$ and Q^s is the characteristic polynomial of the Frobenius action on T_H. Put $T_\ell = \oplus_j T_\ell^{(j)} \oplus T_H$.

Let $m = \nu_\ell(P(1))$, and let $\lambda_1 \le \lambda_2$ be slopes of $\mathrm{Np}_\ell(P(1-t))$. We prove by induction on $\deg f$ that if T is an R-module and the characteristic polynomial of the action of t is $f = P^r Q^s$, then

$$T/(1-x)T \cong \oplus_{j=1}^r G_\ell^{(j)} \oplus (\mathbb{Z}_\ell/Q(1)\mathbb{Z}_\ell)^s,$$

where $\mathrm{Hp}_\ell(G_\ell^{(j)}, 2) = (m_j, m_{2r+s-j})$ such that $m_j \le \lambda_1$, and $m_j + m_{2r+s-j} = m$ for $1 \le j \le r$.

The statement is clear, if $\deg f \le 2$. Assume $\deg f > 2$. Let $\mathrm{Hp}_\ell(T) = (m_1, \dots, m_{2r+s})$. Then

$$T/(1-x)T \cong \oplus_{i=1}^{2r+s} \mathbb{Z}/\ell^{m_i}\mathbb{Z}.$$

Choose $v_1, \dots, v_{2r+s} \in T/(1-x)T$ such that v_i generates the summand $\mathbb{Z}/\ell^{m_i}\mathbb{Z}$. Suppose that $w \in T$ lifts v_{2r+s}. Let $T_1 \subset T$ be the saturated sublattice containing w and Fw. Clearly, T_1 is F-invariant. If the characteristic polynomial of the Frobenius action on T_1 is Q, then $T_1/(1-x)T_1 \cong \mathbb{Z}_\ell/Q(1)\mathbb{Z}_\ell$. Thus $\mathrm{Hp}_\ell(T/T_1) = (m_1, \dots, m_{2r+s-1})$, and we use induction hypothesis.

Suppose that the characteristic polynomial of the Frobenius action on T_1 is P. Then

$$T_1/(1-x)T_1 \cong \mathbb{Z}/\ell^{m-m_{2r+s}}\mathbb{Z} \oplus \mathbb{Z}/\ell^{m_{2r+s}}\mathbb{Z}.$$

By Theorem 3.2,

$$m_1 \le m - m_{2r+s}. \tag{4.1}$$

Now we use the argument from the proof of [Ry12, Theorem 3.4]. By Proposition 2.2, T corresponds to a matrix factorization (X, Y) such that $\det X = f$ and

$$X \equiv \mathrm{diag}(\ell^{m_1}, \dots, \ell^{m_{2r+s}}) \mod (1-t)\mathbb{Z}_\ell[t].$$

The matrix factorization (Y, X) corresponds to a module T' over R, which is generated by $2r+s$ elements and the characteristic polynomial of x is equal to

$$\det Y = P^{2r+s}/f.$$

Moreover,

$$Y \equiv \mathrm{diag}(\ell^{m-m_1}, \dots, \ell^{m-m_{2r+s}}) \mod (1-t)\mathbb{Z}_\ell[t],$$

i.e.,

$$\mathrm{Hp}_\ell(T') = (m - m_{2r+s}, \dots, m - m_1).$$

As before, either $m_1 = \nu_\ell(Q(1))$, and we are reduced to the case $\deg f = r+s-1$, or we have the inequality (4.1) for T':

$$m - m_{2r+s} \le m - (m - m_1) = m_1.$$

It follows that $m = m_1 + m_{2r+s}$.

We proved that

$$T_1/(1-x)T_1 \cong \mathbb{Z}/\ell^{m_1}\mathbb{Z} \oplus \mathbb{Z}/\ell^{m_{2r+s}}\mathbb{Z},$$

thus, by Theorem 3.4, $m_1 \le \lambda_1$.

As in the proof of Theorem 3.2, we have an exact sequence of R-modules:

$$0 \to T_1 \to T \to T_2 = T/T_1 \to 0,$$

and the corresponding sequence of abelian groups:

$$0 \to T_1/(1-x)T_1 \to T/(1-x)T \to T_2/(1-x)T_2 \to 0.$$

Clearly, T_2 is a free $\mathbb{Z}_\ell$-module, and the characteristic polynomial of x on T_2 is $P^{r-1}Q^s$. We claim that

$$\mathrm{Hp}_\ell(T_2) = (m_2, \dots, m_{2r-2+s}).$$

Indeed, $T/(1-x)T$ is generated by $v_1, \dots, v_{2r+s}$ and, by construction of T_1, the group $T_1/(1-x)T_1$ is generated by v_{2r+s} and some v such that $\ell^{m_1}v = 0$. Moreover, since T_2 is generated by $2r+s-2$ elements, we may assume that

$$v = \sum_{j=1}^{2r+s} a_j v_j,\ a_j \in \mathbb{Z}$$

where $\ell^{m_1} a_j v_j = 0$, and there exists j such that $\nu_\ell(a_j) = 0$. In particular, $\nu_\ell(a_j) \geq m_j - m_1$. Thus we can reorder $v_1, \dots, v_j$ and assume that $\nu_\ell(a_1) = 0$. Now we see that the images of $v_2, \dots, v_{2r-2+s}$ generate the subgroup

$$\oplus_{j=2}^{2r-2+s} \mathbb{Z}/\ell^{m_j}\mathbb{Z}$$

of the same order as $T_2/(1-x)T_2$. This proves the claim, and the induction hypothesis for T_2 proves the theorem. □

Theorem 4.2. *Let $f(t) = P(t)(t \pm \sqrt{q})^r$ be a Weil polynomial, where P is separable of degree 2, and $P(\mp\sqrt{q}) \neq 0$. Let*

- *G be a finite abelian group of order $f(1)$;*
- *$\mathrm{Hp}_\ell(G_\ell, r+2) = (m_1, \dots, m_{r+2})$;*
- *$\lambda_1 \leq \lambda_2$ be slopes of $\mathrm{Np}_\ell(P(1-t))$;*
- *$\lambda_q = \nu_\ell(1 \mp \sqrt{q})$.*

Then G is the group of points on some abelian variety with Weil polynomial f if and only if

1. *$\mathrm{Np}_\ell(f(1-t))$ lies on or above $\mathrm{Hp}_\ell(G_\ell, r+2)$;*
2. *$m_r \leq \lambda_q$;*
3. *$m_3 \geq \lambda_q$;*
4. *$m_1 + m_{r+1} \leq \lambda_q + \lambda_1$.*

Proof. Let $G = A(k)$ for some A with Weil polynomial f. Assertion (1) follows from [Ry10, Theorem 1.1], and inequalities (2) and (3) follow from Theorem 3.4 applied to

$$T = T_\ell^{(1)}(A),\ s = r,\ \alpha = 1 \pm \sqrt{q},\ \text{and } \lambda = \lambda_q.$$

In particular, $m_3 = \dots = m_r = \lambda_q$. To prove (4) apply Corollary 3.5 to

$$s = r,\ \alpha = (1 \mp \sqrt{q})^{r-1}(1-\alpha_1) \text{ and } \lambda = (r-1)\lambda_q + \lambda_1,$$

where α_1 is a root of P such that $\nu_\ell(1-\alpha_1) = \lambda_1$. We get that

$$m_1 + m_{r+1} + (r-2)\lambda_q = m_1 + m_3 + \dots + m_{r+1} \leq (r-1)\lambda_q + \lceil \lambda_1 \rceil.$$

Let us prove that if λ_1 is not integral, then

$$m_1 + m_{r+1} < \lambda_q + \lceil \lambda_1 \rceil.$$

In this case $\lambda_1 = \lambda_2 = \nu_\ell(P(1))/2$. If $m_1 + m_{r+1} = \lambda_q + \lceil \lambda_1 \rceil$, then $m_2 + m_{r+2} = \lambda_q + \lfloor \lambda_2 \rfloor$. But from $m_1 + m_{r+1} \leq m_2 + m_{r+2}$ it follows that $\lceil \lambda_1 \rceil \leq \lfloor \lambda_2 \rfloor$.

We prove the theorem in other direction. We construct Tate modules T_ℓ such that $f(t) = \det(t - x|T_\ell)$, and $G_\ell \cong T_\ell/(1-x)T_\ell$ and use Proposition 2.1. If $f \neq f_\ell$ we can apply the construction of the Tate module from the proof of Theorem 4.1. If $\ell = p$, then $\deg P_\ell \leq 1$, and $f \neq f_\ell$. So assume that $f = f_\ell$, and $\ell \neq p$. Let T_1 be a Tate module such that x acts with characteristic polynomial P. Put $T_2 = \mathbb{Z}_\ell$ such that x acts as multiplication by $\mp\sqrt{q}$, and put

$$V = (T_1 \oplus T_2^2) \otimes \mathbb{Q}_\ell.$$

In [Ry12] we construct a Tate module $T \subset V$ such that $\mathrm{Hp}_\ell(T) = (m_1, m_2, m_{r+1}, m_{r+2})$. Put $T_\ell = T \oplus T_2^{r-2}$. □

5. Abelian surfaces and threefolds

By Theorems [Ry10, Theorem 1.1], 4.1 and 4.2 we get a more conceptual proof of [Ry12, Theorem 2.4].

Theorem 5.1. *Let A be an abelian surface over a finite field with Weil polynomial f_A. Let G be an abelian group of order $f_A(1)$. Then G is the group of points on some variety in the isogeny class of A if and only if for any prime number ℓ one of the following conditions holds.*

1. *Suppose f_A has no multiple roots, then $\mathrm{Np}_\ell(f_A(1-t))$ lies on or above $\mathrm{Hp}_\ell(G_\ell, 4)$.*
2. *Suppose $f_A = P_A^2$, and P_A has no multiple roots, then $G_\ell \cong G_\ell^{(1)} \oplus G_\ell^{(2)}$, where $G_\ell^{(1)}$ and $G_\ell^{(2)}$ are ℓ-primary abelian groups with one or two generators such that $\mathrm{Np}_\ell(P_A(1-t))$ lies on or above $\mathrm{Hp}_\ell(G_\ell^{(1)}, 2)$ and $\mathrm{Hp}_\ell(G_\ell^{(2)}, 2)$.*
3. *Suppose $f_A = P(t)(t \pm \sqrt{q})^2$, and $\sqrt{q} \in \mathbb{Z}$, where $P(t)$ is separable, and $P(\mp\sqrt{q}) \neq 0$. Let*
   - *$\mathrm{Hp}_\ell(G_\ell, 4) = (m_1, m_2, m_3, m_4)$;*
   - *$\lambda_1 \leq \lambda_2$ be slopes of $\mathrm{Np}_\ell(P(1-t))$;*
   - *$\lambda_q = \nu_\ell(1 \pm \sqrt{q})$;*

   Then
   (a) *$\mathrm{Np}_\ell(f(1-t))$ lies on or above $\mathrm{Hp}_\ell(G_\ell, 4)$;*
   (b) *$m_2 \leq \lambda_q$;*
   (c) *$m_3 \geq \lambda_q$;*
   (d) *$m_1 + m_3 \leq \lambda_q + \lambda_1$.*
4. *Finally, if $f_A(t) = (t \pm \sqrt{q})^4$, and $\sqrt{q} \in \mathbb{Z}$, then $G \cong (\mathbb{Z}/(1 \pm \sqrt{q})\mathbb{Z})^4$.*

Finally, we give a partial classification of groups of points on abelian threefolds. Note, that in the theorem below we do not know the classification in case (2) for $r > 3$, and in cases (3) and (4) for $r > 2$.

Theorem 5.2. *Let A be an abelian threefold over a finite field with Weil polynomial f_A. Then the following conditions hold for $G = A(k)$:*

1. *Suppose f_A is separable. Then $\mathrm{Np}_\ell(f_A(1-t))$ lies on or above $\mathrm{Hp}_\ell(G_\ell, 6)$.*
2. *Suppose $f_A = P^2$, and P is separable. Let $m = \nu_\ell(P(1))$, and let G_ℓ be generated by r elements. Then $r \geq 2$. If $r = 2$, then $G_\ell \cong (\mathbb{Z}/m\mathbb{Z})^2$, and if $r = 3$, then $\mathrm{Hp}_\ell(G_\ell, 3) = (m_1, m_2, m_3)$, where $\mathrm{Np}_\ell(P(1-t))$ lies on or above the Hodge polygon $(m-m_3, m-m_2, m-m_1)$.*
3. *Suppose $f_A = P^2Q$, where $\deg P = \deg Q = 2$, and the polynomial PQ is separable. Let $m = \nu_\ell(P(1)Q(1))$. Then G_ℓ is generated by $r \geq 2$ elements, and if $r = 2$, then $\mathrm{Hp}_\ell(G_\ell, 2) = (m_1, m_2)$, where $\mathrm{Np}_\ell(Q(1-t))$ lies on or above the Hodge polygon $(m - m_2, m - m_1)$.*
4. *Suppose $f_A(t) = P(t)(t \pm \sqrt{q})^2$, and $\sqrt{q} \in \mathbb{Z}$, where $P(t)$ is separable, and $P(\mp\sqrt{q}) \neq 0$. Let $m = \nu_\ell(P(1)(1 \pm \sqrt{q}))$. Then G_ℓ is generated by $r \geq 2$ elements, and if $r = 2$, then $\mathrm{Hp}_\ell(G_\ell, 2) = (m_1, m_2)$, where $\mathrm{Np}_\ell(P(1-t))$ lies on or above the Hodge polygon $(0, 0, m - m_2, m - m_1)$.*
5. *Suppose $f_A(t) = P(t)(t \pm \sqrt{q})^4$, and $\sqrt{q} \in \mathbb{Z}$, where $P(t)$ is separable, and $P(\mp\sqrt{q}) \neq 0$. Let*
   - *$\mathrm{Hp}_\ell(G_\ell, 6) = (m_1, \ldots, m_6)$;*
   - *$\lambda_1 \leq \lambda_2$ be slopes of $\mathrm{Np}_\ell(P(1-t))$;*
   - *$\lambda_q = \nu_\ell(1 \pm \sqrt{q})$.*

   Then
   (a) *$\mathrm{Np}_\ell(f(1-t))$ lies on or above $\mathrm{Hp}_\ell(G_\ell, 6)$;*
   (b) *$m_2 \leq \lambda_q$;*

(c) $m_3 = m_4 = \lambda_q$;
(d) $m_5 \geq \lambda_q$;
(e) $m_1 + m_5 \leq \lambda_q + \lambda_1$.

(6) *Finally, suppose* $f_A = P^r Q^s$ *for some* r *and* s, *where* $\deg P \leq 2$, *the polynomial* P *is separable, and* Q *divides* P. *Then* $G_\ell \cong \oplus_{j=1}^r G_\ell^{(j)} \oplus H$, *where* $G_\ell^{(j)}$ *are* ℓ*-primary abelian groups such that* $\mathrm{Np}_\ell(P(1-t))$ *lies on or above* $\mathrm{Hp}_\ell(G_\ell^{(j)}, 2)$ *for all* $1 \leq j \leq r$, *and* $H \cong (\mathbb{Z}_\ell/Q(1)\mathbb{Z}_\ell)^s$.

Conversely, let G *be an abelian group of order* $f_A(1)$ *such that one of the conditions above holds. Then* G *is the group of points on some variety in the isogeny class of* A.

Proof. The cases (1), (5) and (6) follow from Theorems [Ry10, Theorem 1.1], 4.1 and 4.2. We prove (2) first. Without loss of generality assume that $P = P_\ell$. Recall that $P(x) = 0$, and $\nu_\ell(P(1)) = m$. It follows that the group

$$G_\ell \cong T_\ell(A)/(1-x)T_\ell(A)$$

is annihilated by ℓ^m. Thus $r \geq 2$, and if $r = 2$, then $G_\ell \cong (\mathbb{Z}/m\mathbb{Z})^2$. Assume $r = 3$. By Proposition 2.1, we have to find out when there exists a module T over $R = \mathbb{Z}_\ell[t]/P(t)\mathbb{Z}_\ell[t]$ with $r = 3$ generators such that $\mathrm{Hp}_\ell(T/(1-x)T, 3) = (m_1, m_2, m_3)$. By Proposition 2.2, there exists a matrix factorization (X, Y) such that $\det X = f_A$ and

$$X \equiv \mathrm{diag}(\ell^{m_1}, \ell^{m_2}, \ell^{m_3}) \mod (1-t)\mathbb{Z}_\ell[t].$$

The matrix factorization (Y, X) corresponds to a module T' over R, which is generated by 3 elements, and the characteristic polynomial of x is equal to

$$\det Y = P^3/f_A = P,$$

and

$$Y \equiv \mathrm{diag}(\ell^{m-m_1}, \ell^{m-m_2}, \ell^{m-m_3}) \mod (1-t)\mathbb{Z}_\ell[t].$$

Such a module exists if and only if $\mathrm{Np}_\ell(P(1-t))$ lies on or above $\mathrm{Hp}_\ell(T'/(1-x)T', 3) = (m - m_3, m - m_2, m - m_1)$. We proved that there exists a module T such that $\mathrm{Hp}_\ell(T/(1-x)T, 3) = (m_1, m_2, m_3)$ if and only if $\mathrm{Np}_\ell(P(1-t))$ lies on or above $(m - m_3, m - m_2, m - m_1)$.

In cases (3) and (4) the group G_ℓ is annihilated by ℓ^m. Thus it is not cyclic, and $r \geq 2$. We now consider the case (3). Let $R = \mathbb{Z}_\ell[t]/P(t)Q(t)\mathbb{Z}_\ell[t]$. Assume we have an R-module T with $r = 2$ generators such that $\mathrm{Hp}_\ell(T/(1-x)T, 2) = (m_1, m_2)$. As before, there exists a matrix factorization (X, Y) such that $\det X = f_A$ and

$$X \equiv \mathrm{diag}(\ell^{m_1}, \ell^{m_2}) \mod (1-t)\mathbb{Z}_\ell[t].$$

The matrix factorization (Y, X) corresponds to a module T' over R, which is generated by 2 elements, and the characteristic polynomial of x is equal to

$$\det Y = (PQ)^2/f_A = Q,$$

and

$$Y \equiv \mathrm{diag}(\ell^{m-m_1}, \ell^{m-m_2}) \mod (1-t)\mathbb{Z}_\ell[t].$$

As above, it follows that there exists a module T such that $\mathrm{Hp}_\ell(T/(1-x)T, 2) = (m_1, m_2)$ if and only if $\mathrm{Np}_\ell(Q(1-t))$ lies on or above $\mathrm{Hp}_\ell(T'/(1-x)T', 3) = (m - m_2, m - m_1)$.

Case (4). Let $R = \mathbb{Z}_\ell[t]/P(t)(t \pm \sqrt{q})\mathbb{Z}_\ell[t]$, and let T be an R-module with $r = 2$ generators such that $\mathrm{Hp}_\ell(T/(1-x)T, 2) = (m_1, m_2)$. For the corresponding matrix factorization (X, Y) the matrix factorization (Y, X) gives a module T' over R, which is generated by 2 elements, and the characteristic polynomial of x is equal to

$$\det Y = (P(t \pm \sqrt{q}))^2/f_A = P,$$

and

$$Y \equiv \mathrm{diag}(\ell^{m-m_1}, \ell^{m-m_2}) \mod (1-t)\mathbb{Z}_\ell[t].$$

The rank of T' is 4, and such a module exists if and only if $\mathrm{Np}_\ell(P(1-t))$ lies on or above $\mathrm{Hp}_\ell(T'/(1-x)T', 4) = (0, 0, m-m_2, m-m_1)$. Therefore there exists a module T such that $\mathrm{Hp}_\ell(T/(1-x)T, 2) = (m_1, m_2)$ if and only if $\mathrm{Np}_\ell(P(1-t))$ lies on or above $\mathrm{Hp}_\ell(T'/(1-x)T', 3) = (m-m_2, m-m_1)$. □

References

[La94] S. Lang. *Algebraic number theory.* Graduate Texts in Mathematics, Springer, 1994.

[Ry10] S. Rybakov. *The groups of points on abelian varieties over finite fields.* Cent. Eur. J. Math. 8(2), 2010, 282-288. arXiv:0903.0106

[Ry12] S. Rybakov. *The groups of points on abelian surfaces over finite fields.* In Arithmetic, Geometry, Cryptography and Coding Theory, Cont. Math., vol. 574, Amer. Math. Soc., Providence, RI, 2012, pp. 151-158. arXiv:1007.0115

[Ry14] S. Rybakov. *The finite group subschemes of abelian varieties over finite fields.* Finite Fields and Their Applications. 29 (2014), 132-150. arXiv:1006.5959

[Ta66] J. Tate. *Endomorphisms of abelian varieties over finite fields.* Inventiones mathematicae 1966, Volume 2, Issue 2, pp 134–144.

[Wa69] Waterhouse W., Abelian varieties over finite fields, Ann. scient. Éc. Norm. Sup., 1969, 4 serie 2, 521–560.

[WM69] Waterhouse W., Milne J., Abelian varieties over finite fields, Proc. Sympos. Pure Math., Vol. XX, State Univ. New York, Stony Brook, N.Y., 1969, 53–64.

Institute for information transmission problems of the Russian Academy of Sciences

Poncelet laboratory (UMI 2615 of CNRS and Independent University of Moscow)

AG Laboratory, HSE, 7 Vavilova str., Moscow, Russia, 117312

E-mail address: rybakov@mccme.ru, rybakov.sergey@gmail.com